



TEST SAVOLLARIDAN NAMUNALAR

1. b soni $f(x)$ funksiyaning $x \rightarrow +\infty$ dagi limiti deyiladi, agar har qanday $\varepsilon > 0$ son uchun shunday:

- A) $(M; +\infty)$ oraliq topilsaki, unda $f(x) - b < \varepsilon$ tengsizlik bajarilsa;
- B) $(M; +\infty)$ oraliq topilsaki, unda $|f(x) - b| < \varepsilon$ tengsizlik bajarilsa;
- D) $(M; -\infty)$ oraliq topilsaki, unda $|f(x) - b| < \varepsilon$ tengsizlik bajarilsa;
- E) $(M; -\infty)$ oraliq topilsaki, unda $f(x) - b < \varepsilon$ tengsizlik bajarilsa;
- F) $(M; +\infty)$ oraliq topilsaki, unda $|f(x) - b| > \varepsilon$ tengsizlik bajarilsa.

2. Agar $f(x)$ va $\varphi(x)$ funksiyalar $x \rightarrow a$ da mos ravishda b va c ga teng limitlarga ega bo'lsa, ularning $f(x) + g(x)$ yig'indisi

- A) $x \rightarrow +\infty$ da $b + c$ limitga ega bo'ladi;
- B) $x \rightarrow -\infty$ da $b + c$ limitga ega bo'ladi;
- D) $x \rightarrow \infty$ da $b + c$ limitga ega bo'ladi;
- E) $x \rightarrow a$ da $b + c$ limitga ega bo'ladi;
- F) $x \rightarrow a$ da $|b + c|$ limitga ega bo'ladi.

3. $f(x)$ funksiya $x = a$ nuqtada uzluksiz deyiladi,

- A) agar $f(x) = f(a)$ bo'lsa;
- B) agar $f(a - 0) \neq f(a + 0)$ bo'lsa;
- D) agar $\lim_{x \rightarrow a} f(x) = \infty$ bo'lsa;
- E) agar $\lim_{x \rightarrow a} f(x) = -f(x)$ bo'lsa;
- F) agar $f(x)$ funksiya $x = a$ nuqtada aniqlangan va $f(x) - f(a)$ ayirma $x \rightarrow a$ da cheksiz kichik bo'lsa.

4. Agar $f(x)$ va $g(x)$ funksiyalar $x = a$ nuqtada aniqlangan bo'lsa, u holda

- A) ularning faqat yig'indisi (ayirmasi va ko'paytmasi emas) shu nuqtada uzluksiz bo'lishi mumkin;
- B) ularning yig'indisi va ayirmasi (ko'paytmasi emas) shu nuqtada uzluksiz bo'lishi mumkin;
- D) ularning yig'indisi, ayirmasi, ko'paytmasi ham shu nuqtada uzluksiz bo'ladi;
- E) ularning yig'indisi, ayirmasi, ko'paytmasi ham shu nuqtada uzluksiz bo'lishi mumkin;
- F) ularning yig'indisi, ayirmasi, ko'paytmasi shu nuqta yotgan oraliqda uzluksiz bo'ladi.

5. Agar $f(x)$ va $g(x)$ funksiyalar $x = a$ nuqtada uzluksiz bo'lsa, u holda

- A) $\frac{f(x)}{g(x)}$ funksiya ham shu nuqtada uzluksiz bo'ladi;
- B) $\frac{f(x)}{g(x)}$ va $\frac{g(x)}{f(x)}$ funksiyalar ham shu nuqtada uzluksiz bo'ladi;
- D) $\frac{1}{f(x)} \cdot \frac{1}{g(x)}$ funksiya ham shu nuqtada uzluksiz bo'ladi;
- E) $g(x) \neq 0$ bo'lganda $\frac{f(x)}{g(x)}$ funksiya ham shu nuqtada uzluksiz bo'ladi;
- F) $f(x) \neq 0$ bo'lganda $\frac{f(x)}{g(x)}$ funksiya ham shu nuqtada uzluksiz bo'ladi.

6. Oraliqning (intervalning) barcha nuqtalarida uzluksiz bo'lgan funksiya

- A) shu oraliqning (intervalning) ayrim nuqtalarida uzluksiz deyiladi;
- B) shu oraliqning (intervalning) faqat nuqtalarida uzluksiz deyiladi;
- D) shu oraliqning (intervalning) faqat o'rtasida uzluksiz deyiladi;
- E) shu oraliqda (intervalda) uzluksiz deyiladi;
- F) shu oraliqning faqat ko'rsatilgan qismida uzluksiz bo'ladi, deyiladi.

7. Agar x radianlarda berilgan bo'lsa, u holda $\lim_{x \rightarrow 0} \frac{\sin x}{x} =$

- A) 0; B) 1; D) -1; E) $-\infty$; F) $+\infty$.

8. Agar x radianlarda berilgan bo'lsa, u holda $\lim_{x \rightarrow a} \sin x =$

- A) a ; B) $\sin a$; D) $\frac{\sin x}{a}$; E) $a \sin x$; F) $\sin \frac{x}{2}$.

9. Agar $\alpha(x)$ o'zgarmas funksiya $x \rightarrow +\infty$ da cheksiz kichik bo'lsa,

- A) x ning barcha qiymatlarida $\alpha(x) = 0$ bo'ladi;
- B) x ning barcha qiymatlari $\alpha(x) = +\infty$ bo'ladi;
- D) $x = 0$ da $\alpha(x) = 0$ bo'ladi;
- E) $x = 0$ da $\alpha(x) = -\infty$ bo'ladi;
- F) $x = 0$ da $\alpha(x) = 1$ bo'ladi.

10. Agar $P(x) = a_m x^m + a_{m-1} x^{m-1} + \dots + a_0$ va $Q(x) = b_n x^n + b_{n-1} x^{n-1} + \dots + b_0$, $a_m \neq 0$, $b_n \neq 0$ va ko'phadlarning darajalari $m < n$

bo'lsa, u holda $\lim_{x \rightarrow \infty} \frac{P(x)}{Q(x)} = \dots$ bo'ladi.

- A) $\frac{m}{n}$; B) $\frac{a_0}{b_0}$; D) $\frac{a_m}{b_n}$; E) 0; F) ∞ .

11. a ning h radiusli teshilgan (o'yilgan) atrofi

- A) shu nuqtaning o'zi chiqarib tashlangan atrofidan iborat;
 B) $(a - h; a)$ va $(a; a + h)$ oraliqlarning birlashmasidan iborat;
 D) $(-\infty; a)$ va $(a; +\infty)$ oraliqlarning birlashmasidan iborat;
 E) $(-\infty; h)$ va $(h; +\infty)$ oraliqlarning birlashmasidan iborat;
 F) $(-h; a)$ va $(a; h)$ oraliqlarning birlashmasidan iborat.

12. Agar $[a; b]$ yarim intervalda berilgan $f(x)$ funksiya uchun $\lim_{x \rightarrow b-0} f(x) = +\infty$ bo'lsa, $x = b$ to'g'ri chiziq $f(x)$ funksiya grafigi uchun:

- A) gorizontal asimptota; B) vertikal asimptota;
 D) gorizontal urinma; E) vertikal urinma;
 F) og'ma asimptota.

13. Agar f funksiya $[a; b]$ kesmada o'suvchi (kamayuvchi) va uzluksiz bo'lsa, u holda shu funksiyaga

- A) $[a; b]$ kesmada (mos ravishda $[b; a]$ kesmada) aniqlangan f^{-1} teskari funksiya mavjud;
 B) $[a; b]$ kesmada aniqlangan f^{-1} teskari funksiya mavjud;
 D) $[f(a); f(b)]$ kesmada (mos ravishda $[f(b); f(a)]$ kesmada) aniqlangan f^{-1} teskari funksiya mavjud bo'ladi;
 E) $[f(b); f(a)]$ kesmada (mos ravishda $[f(a); f(b)]$ kesmada) aniqlangan f^{-1} teskari funksiya mavjud;
 F) $[f(a); f(b)]$ kesmada aniqlangan f^{-1} teskari funksiya mavjud.

14. $\lim_{x \rightarrow 1} \frac{x^6 - 1}{x^3 - 1} = \dots$

- A) 0; B) 2; D) 3; E) $+\infty$; F) $-\infty$.

15. $f(x) = kx + b$ to'g'ri chiziq f funksiya grafigining $x \rightarrow \infty$ dagi og'ma asimptotasi bo'lishi uchun ... bo'lishi zarur va yetarli.

- A) $k = \lim_{x \rightarrow \infty} f(x)$, $b = \lim_{x \rightarrow \infty} (f(x) - k)$;
 B) $k = x$, $b = \lim_{x \rightarrow \infty} f(x)$;

$$D) k = \lim_{x \rightarrow \infty} (f(x) - b), b = \lim_{x \rightarrow \infty} (f(x) - kx);$$

$$D) k = \lim_{x \rightarrow \infty} \left(\frac{f(x)}{x} - b \right), b = \lim_{x \rightarrow \infty} \frac{f(x)}{x};$$

$$E) k = \lim_{x \rightarrow \infty} \frac{f(x)}{x}, b = \lim_{x \rightarrow \infty} (f(x) - kx).$$

16. $f(x)$ funksiyaning $x = a$ nuqtada chapdan (shu kabi o'ngdan) uzluksiz bo'lishi uchun ... bo'lishi zarur.

A) $f(a - 0) = f(0)$ (mos ravishda $f(a + 0) = f(0)$);

B) $f(a - 0) = 0$ (mos ravishda $f(a + 0) = 0$);

C) $f(a - 0) \neq f(a + 0)$;

D) $f(a - 0) = f(a + 0)$;

E) $f(a - 0) = f(a)$ (mos ravishda $f(a + 0) = f(a)$).

17. Agar $f(x)$ funksiya $[a; b]$ kesmada uzluksiz, monoton va $f(a)f(b) < 0$ bo'lsa, funksiya shu oraliqning ... nuqtasida nolga aylanadi.

A) faqat bir;

B) tasodifan bir;

D) hech bir nuqtasida nolga aylanmaydi;

E) $f(a)f(b) > 0$ bo'lsa, bir;

F) kamida bir.

18. $y = f(x)$ funksiyada $x = x_0$ nuqtada olingan $f'(x_0)$ hosila deb

A) har qanday $\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$ limitga aytiladi, bunda $\Delta y = f(x + \Delta x) - f(x)$ funksiya orttirmasi, Δx - argument orttirmasi;

B) $\frac{\Delta y}{\Delta x}$ nisbatga aytiladi, bunda Δy - funksiya orttirmasi, Δx - argument orttirmasi;

D) chekli $\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}$ limitga aytiladi, bunda Δy - funksiya orttirmasi, Δx - argument orttirmasi;

E) chekli $\lim_{x \rightarrow \infty} (\Delta y - \Delta x)$ limitga aytiladi, bunda Δy - funksiya orttirmasi, Δx - argument orttirmasi;

F) $\lim_{x \rightarrow \infty} \frac{\Delta y}{\Delta x}$ limitga aytiladi, bunda Δy - funksiya orttirmasi, Δx - argument orttirmasi.

19. Agar biror y kattalik $y = f(x)$ qonun bo'yicha o'zgarayotgan bo'lsa, bu kattalikning $x = x_0$ dagi o'zgarish oniy tezligi ... ga teng.

A) $f(x_0)$; B) $\frac{\Delta f(x_0)}{\Delta x_0}$; D) $f'(x_0)$; E) $f(x_0 + \Delta x) - f(x_0)$; F) $\frac{f(x_0)}{x_0}$.

20. $A(x_0; y_0)$ nuqtada $y=f(x)$ egri chiziqqa o'tkazilgan urinmaning k burchak koeffitsiyenti ... ga teng.

A) $\frac{f(x_0)}{x_0}$; B) $f(x_0 + \Delta x) - f(x_0)$; D) $f(x_0)$; E) $f'(x_0)$; F) $\frac{\Delta f(x_0)}{\Delta x_0}$.

21. Agar $f(x)$ va $g(x)$ funksiyalar $f'(x)$, $g'(x)$ hosilalari mavjud bo'lsa, u holda $(f(x) \pm g(x))' =$

A) $f'(x) \cdot g'(x)$; B) $f'(x) \pm g'(x)$; D) $f'(x \pm y)$; E) $g'(x \pm y)$; F) $f(x) \pm g(x)$.

22. Agar $f'(x)$ va $g'(x)$ hosilalar mavjud bo'lsa, $(f(x)g(x))' =$

A) $f'(x)g'(x)$; B) $f'(x)g'(x) + C$, C - o'zgarmas;
 D) $f'(x)f(x) + g'(x)g(x)$; E) $f'(x)g(x) + g'(x)f(x)$;
 F) $f'(x)g'(x) + f(x)g(x)$.

23. Agar $f'(x)$ va $g'(x)$ hosilalar mavjud va $g(x) \neq 0$ bo'lsa, $\left(\frac{f(x)}{g(x)}\right)' =$

A) $\frac{f'(x)}{g'(x)}$; B) $\frac{f'(x)}{g(x)}$;
 D) $\frac{f'(x)g(x) - f(x)g'(x)}{g^2(x)}$; E) $\frac{f'(x)g(x) - f(x)g'(x)}{g(x)}$;
 F) $\frac{f'(x)g(x) + f(x)g'(x)}{g^2(x)}$.

24. Berilgan $f(x) = x^a$, $a \in R$ funksiyaning $f'(x)$ hosilasi ifodasini ko'rsating:

A) $a \cdot x^a$; B) 1; D) 0; E) $a \cdot x^{a-1}$; F) $f''(x)$.

25. $f(x)$ funksiyalarning $f'(x)$ hosilalari ifodasini ko'rsating:

$f(x) =$	$f'(x) =$
1) $\sin x$; 2) $\cos x$;	K) $\sin x$; L) $-\sin x$; M) $\cos x$;
3) $\operatorname{tg} x$; 4) $\operatorname{ctg} x$;	N) $-\frac{1}{\cos^2 x}$; P) $\frac{1}{\cos^2 x}$; Q) $-\frac{1}{\sin^2 x}$.

A) 1K, 4P, 3M, 2N; B) 1M, 4Q, 3P, 2L;
 D) 2L, 4K, 3Q, 1N; E) 3R, 4K, 2M, 1Q;
 F) 4Q, 3N, 2K, 1L.

26. $f(x)$ funksiyalarning $f'(x)$ hosilalari ifodasini ko'rsating:

$f(x) =$	$f'(x) =$
1) e^x ; 2) a^x ($a > 0$, $a \neq 1$);	K) $\frac{1}{x \ln a}$; L) e^x ; M) $\ln a^x$;

32. Differensiallanuvchi funksiya $x = c$ nuqtada ekstremumga ega bo'lishi uchun $f'(c) = 0$ bo'lishi

- A) yetarli; B) zarur va yetarli; D) zarur;
E) yetarli, lekin zaruriy emas; F) shart emas.

33. Funksiya hosilasi mavjud bo'lmagan nuqtada funksiya ekstremumga ...

- A) ega bo'lmaydi; B) har vaqt ega bo'ladi;
D) faqat minimumga ega bo'ladi; E) ega bo'lishi mumkin;
F) faqat maksimumga ega bo'ladi.

34. $f(x)$ funksiya $c \in (a; b)$ nuqtada hosilaga ega bo'lsin. Agar $f'(c) = 0$ va c nuqtadan chapda $f' > 0$, nuqtadan o'ng tomonda $f' < 0$ bo'lsa, funksiya $x = c$ nuqtada ... ga erishadi.

- A) lokal minimum;
B) maksimum yoki minimum ;
D) lokal maksimum;
E) intervaldagi eng kichik qiymat;
F) intervaldagi eng katta qiymat.

35. $f''(x)$ hosila $x = c$ nuqtada 0ga teng. Bu nuqta $f(x)$ funksiya uchun qanday nuqtadan iborat?

- A) minimum; B) bukilish; D) maksimum;
E) ekstremum; F) uzilish.

36. Lagranj teoremasi: agar $f(x)$ funksiya $[a; b]$ kesmada uzluksiz va oraliqning ichki nuqtalarida differensiallansa, bu kesmada shunday $x = s$ nuqta topiladiki, unda ... tenglik o'rinli bo'ladi.

- A) $\frac{f(b)-f(a)}{b-a} = f'(c)$; B) $\frac{f(b)+f(a)}{b+a} = f'(c)$;
C) $(f(b) - f(a))(b - a) = f'(c)$; D) $\frac{f(x)-f(b)}{f(b)-f(a)} = f'(c)$;
E) $\frac{f(x)+f(b)}{x+a} = f'(c)$.

37. Agar $[a; b]$ kesmada $f(x)$ funksiya uzluksiz va $f'' > 0$ bo'lsa, funksiya grafigi qavariqligi bilan ... qaragan bo'ladi.

- A) yuqoriga; B) o'ngga; D) pastga; E) har tomonga; F) chapga.

38. $(x+a)^n$ Nyuton binomi yoyilmasidagi $(k+1)$ - hadi ... ko'rinishda bo'ladi.

- A) $C_n^k x^n a^k$; B) $C_{n-k}^k x^{n-k} a^n$; D) $C_{n+k}^k x^k a^{n-k}$;
 E) $C_n^k x^{n-k} a^k$; F) $C_{n+k}^{n-k} x^n a^k$.

39. Har qaysi $f(x)$ funksiyaning $F(x)$ boshlang'ich funksiyasini ko'rsating:

$f(x) =$		$F(x) =$
A) $\sqrt{3x-2}$; B) $\frac{2}{\sqrt{3x-2}}, x > \frac{2}{3}$		K) $\frac{3x^2}{\sqrt{3x-2}}$; L) $\frac{2}{9}\sqrt{(3x-2)^3}$; P) $\frac{2}{3}\sqrt{3x-2}$; N) $(3x-2)\sqrt{3x-2}$;

- 1) AK, BL; 2) AP, BK; 3) AL, BP; 4) AN, BP; 5) AL, BN.

40. $\int f(x)dx = F(x) + C, C \in R$ bo'yicha har qaysi $f(x)$ funksiyaga qaysi $F(x)$ funksiya mos?

$f(x) =$		$F(x) =$
K) $\frac{1}{1+x^2}$; L) $\cos x$; M) $\frac{1}{\sin^2 x}$		A) $\frac{(1+x^2)^2}{2}$; B) $\sin x$; D) $-\sin x$; E) $-\operatorname{tg} x$; F) $-\operatorname{ctg} x$; G) $\operatorname{arctg} x$; H) $\operatorname{arcsin} x$.

- 1) KA, LD, ME; 2) KG, LB, MF; 3) KA, LF, MG;
 4) KE, LG, MH; 5) KD, LB, MB.

41. $\int \left(2x^2 - \frac{1}{x^2} \right) dx$ integralni toping:

- A) $x^3 - \operatorname{arctg}(x-1) + C$; B) $\frac{2x^2 - \frac{1}{x^2}}{2} + C$;
 D) $2x^3 - \ln x^2 + C$; E) $\frac{2x^3}{3} + \frac{1}{x} + C$;
 F) $2x^3 - 2 \ln x + C$.

42. $\int \frac{10 \sin^2 x - 4 \cos^2 x}{\sin^2 x \cos^2 x} dx$ integralni hisoblang.

- A) $10 \operatorname{tg} x + 4 \operatorname{ctg} x + C$; B) $6 \operatorname{tg} x - 8 \sin 2x + C$;
 D) $-\frac{10}{\cos x} + \frac{1}{\sin x} + C$; E) $10 \ln(\cos^2 x) - \frac{1}{\sin x} + C$;
 F) $6 \operatorname{ctg} x - 8 \cos 2x + C$.

43. O'zgaruvchini almashtirishdan foydalanib hisoblang:

$$J = \int \frac{\operatorname{ctg}^3 4x dx}{\sin^2 4x} \text{ va } K = \int \frac{dx}{x \ln x}.$$

- A) $J = \frac{\operatorname{tg}^4 4x}{16}$, $K = x^{\ln x}$; | B) $J = -\frac{\operatorname{ctg} 4x}{16} + C$, $K = \ln(\ln x) + C$;
 D) $J = \cos 4x$, $K = \frac{1}{\ln x}$; | E) $J = \sin 4x$, $K = e^{-x}$;
 F) $J = \operatorname{tg} 4x$, $K = \ln^2 x$.

44. O'zgaruvchilarni ajratishdan foydalanib, $y' = x^3 y^3$ differensial tenglamaning $y(1) = -4$ boshlang'ich shartni qanoatlantiruvchi yechimini toping.

- A) $16x^3$; | B) $-4x^4$; | D) $-4x^{-4}$; | E) $-16x^3$; | F) $-16x^{-3}$.

45. $S_1 = \int_0^2 x^2 \sqrt{1+x^2} dx - ?$ $S_2 = \int_0^2 \frac{xdx}{\sqrt{1-x^4}} - ?$

- A) $S_1 = \frac{9}{52}$, $S_2 = 2 \arcsin 2$; B) $S_1 = 52$, $S_2 = \arccos 4$;
 D) $S_1 = \frac{26}{9}$, $S_2 = 2 \arccos 4$; E) $S_1 = \frac{54}{9}$, $S_2 = \sqrt{1-x^4}$;
 F) $S_1 = \frac{52}{9}$, $S_2 = \frac{1}{2} \arcsin 4$.

46. $\lim_{x \rightarrow +\infty} \left(\frac{3x+15}{3x+1} \right)^{x-1}$ ni hisoblang.

- A) 15; B) $e^{\frac{14}{3}}$; D) $\frac{\infty}{\infty}$; E) e^{15} ; F) 1.

47. Bir aylanada yotgan besh nuqta ustidan qancha vatar o'tkazish mumkin?

- A) C_5^2 ; B) A_5^2 ; D) P_5 ; E) \bar{A}_5^2 ; F) \bar{C}_5^2 .

48. Cheksiz kamayuvchi geometrik progressiyada:

$$S = \frac{3}{4}, \quad a_1 = -\frac{1}{3}; \quad a_n - ?$$

- A) $\frac{5^{n-1}}{2^{2n-1}}$; B) $\frac{9}{56}$; D) $\frac{1}{91}$; E) $\frac{3^2}{5}$; F) 0.

49. $y = x^3 - 3x$ chiziq va uning $x_0 = -1$ absissali nuqtadagi urinmasi bilan chegaralangan shaklning yuzini toping.

A) 5,25; B) 6,75; D) 6,25; E) 4,75; F) 5,75.

50. I integraldan har biri qaysi K ifodaga tengligini va ... nuqtalar o'rnida turgan ifodani ko'rsating.

I: A) $\int_a^b f(x) dx$; B) $\int_a^b f(x) dx$;
 D) $\int_c^d f(x) dx = \dots + \int_e^d f(x) dx$; E) $\int_a^b [f(x) + q(x)] dx$;

K: 1) $-\int_a^b f(x) dx$; 2) $\int_c^e f(x) dx$; 3) $-\int_a^b f(x) dx$; 4) 0;

5) $\int_a^b f(x) dx + \int_a^b q(x) dx$; 6) $\int_a^b f(x) dx - \int_a^b q(x) dx$;

7) $\int_c^{d-e} f(x) dx$; 8) $\int_b^{2b} f(x) dx$; 9) C; 10) $\int_{2a}^{2b} f(x) dx$.

A) A3, B1, D7, E4; B) A1, B4, D2, E5;
 D) A5, B8, D6, E9; E) A10, B9, D4, E6;
 F) A8, B6, D9, E10.

51. Agar $[a; b]$ kesmada $f(x) \geq 0$ funksiya uchun $k \leq f(x) \leq K$

tengsizlik o'rinli bo'lsa, $?(b-a) \leq \int_a^b f(x) dx \leq K?$ bo'ladi. ? belgilar

o'rniga mos ifodalarni tartibi bo'yicha yozing:

A) $(k - a), (k - b)$; B) $(K - k), (K + k)$; D) $k, (b - a)$;
 E) $(k - b), (k + b)$; F) $(K - a), (k + b)$.

52. $f(x)$ funksiya $[a; b]$ kesmada monoton o'suvchi. Agar $[a; b]$ kesma teng n bo'lakka bo'lingan va bo'linish nuqtalari $a = x_0 < x_1 < \dots < x_n = b$ bo'lsa, u holda

$$\frac{?}{n} \sum_{k=0}^{n-1} ? \leq \int_a^b f(x) dx \leq \frac{?}{n} \sum_{k=1}^n ?$$

bo'ladi. ? belgilari o'rniga mos ifodalarni tartibi bilan yozing.

A) $x_n, f(x), x_{n-1}, n - 1, x_k$; B) $x_0, x_k, x_n, n - 2, f(x_{k-1})$;
 D) $b + a, f(x_{k-1}), b + a, n - 1, f(x_{k-1})$; E) $b - a, f(x_k), b - a, n, f(x_k)$;

F) $x_n - x_{n-1}, f(x_{k-2}), x_n - x_{n-1}, 2, f(x_{k-2})$.

53. Trapetsiyalar formulasi:

$$\int_a^b f(x) dx \approx \frac{b-a}{n} \left(\frac{f(a)+f(b)}{2} + f(x_1) + \dots + f(x_{n-1}) \right)$$

? belgilar o'rniga mos ifodalarni kelish tartibida yozing.

A) $b - a, b, x_{n-1};$ B) $b + a, na, x_n;$ D) $ab, x_{n-1}, x_{n-2};$

E) $\sqrt{ab}, b - a, nx_0;$ F) $\sqrt{\frac{b}{a}}, nb, nx.$

54. $x > 0, a > 0$ uchun

Topilsin:	Javob variantlari:				
	(1)	(2)	(3)	(4)	(5)
$(\ln(ax))'$	$\frac{1}{ax}$	$\frac{1}{x}$	$\frac{a}{x}$	$\ln a + \ln x$	e^{ax}
$x = \frac{1}{e^2}$ da $\ln x$	2	-2	e^2	e^{-2}	1
$\lim_{x \rightarrow +\infty} \ln x$	0	$+\infty$	$-\infty$	1	e
$\lim_{x \rightarrow +0} \ln x$	$+\infty$	$-\infty$	-1	e	0

55. $a > 0, b > 0$ uchun:

Topilsin:	Javob variantlari:				
	(1)	(2)	(3)	(4)	(5)
$a^x \cdot a^y$	a^{xy}	$a^x + a^y$	a^{x+y}	a^{x-y}	$a^x \cdot y$
$\frac{a^x}{a^y}$	$a^{\frac{x}{y}}$	$a^x - a^y$	a^{x-y}	$\sqrt[y]{a^x}$	$\frac{a^x}{y}$
$(a^x)^y$	$a^x \cdot y$	a^{x+y}	a^{xy}	$\sqrt[x]{a^y}$	$(a^x)^{a^y}$
$(ab)^x$	$a^x + b^x$	ab^x	$a^x b^x$	$a^x b$	$(a+b)^x$
$\left(\frac{a}{b}\right)^x$	$a^x - b^x$	$\frac{a}{b^x}$	$\frac{a^x}{b^x}$	$\frac{a^x}{b}$	$(a-b)^x$

56. Kombinatorika elementlari:

Asosiy formulalar	Javob variantlari:				
	(1)	(2)	(3)	(4)	(5)
$\bar{A}_m^k =$	$(m!)^k =$	$m \cdot k$	$m! \cdot k!$	k^m	m^k
$A_m^k =$	$\frac{k!}{m!}$	$\frac{m!}{k!}$	$\frac{(m-k)!}{(m+k)!}$	$\frac{(m+k)!}{(m-k)!}$	$\frac{m!}{(m-k)!}$

davomi

$P_m =$	$m!(m-1)!...1!$	$(m-1)!$	$(m+1)!$	$m(m-1)$	$m!$
$C_m^k =$	$\frac{k!}{(m+k)!}$	$\frac{k+1}{m!}$	$\frac{k-1}{m!}$	$\frac{(m+1)!}{m!(m+k)!}$	$\frac{k!}{k!(m-k)!}$
$k=k_1+\dots+k_m$ $P(k_1, k_2, \dots, k_m) =$	$k!$	$\frac{(k+1)!}{k_1!k_2!\dots k_m!}$	$\frac{k}{k_1!k_2!\dots k_m!}$	$\frac{k+1}{k_1!k_2!\dots k_m!}$	$\frac{k!}{k_1!k_2!\dots k_m!}$
$\bar{C}_m^k =$	C_{m+1}^{k+1}	C_{k-m+1}^{k-1}	C_{k+m}^{k-1}	C_{k-m-1}^{k+1}	C_{k+m-1}^k

57. Ehtimollik nazariyasi elementlari:

Qo'shish teoremlari	Javob variantlari:				
	(1)	(2)	(3)	(4)	(5)
$A \cup B = \emptyset$ uchun					
$P(A \cup B) =$	$P(A) \cup P(B)$	$P(A) + P(B)$	$P(A) - P(B)$	$P(A) \cap P(B)$	$P(A - B)$
$P(\bar{A}) =$	$\bar{P(A)}$	$1 - P(A)$	$1 + P(\bar{A})$	$1 - P(\bar{A})$	$1 + P(A)$

58. Bitta ehtimollik fazosidan olingan erkli A va B tasodifiy hodisalar uchun:

	Javob variantlari:				
	(1)	(2)	(3)	(4)	(5)
$P(A \cap B) =$	$P(A) \cdot P(B)$	$P(A) + P(B)$	$P(\bar{A} \cap B)$	$P(A \cap \bar{B})$	$P(\bar{A} \cap \bar{B})$
$P(A \cap B) =$	$P(A) + P(B) - P(A) \cdot P(B)$	$P(A + B) - P(A) \cdot P(B)$	$P(A + B) + P(A) \cdot P(B)$	$P(A) + P(B) + P(A) \cdot P(B)$	$P(A) - P(B) + P(A) \cdot P(A)$

59. X hodisa ro'y bergandagina A hodisaning ro'y berish ehtimolligi $P(A|X) =$

- A) $\frac{P(A \cup X)}{P(X)}$; B) $\frac{P(A \cap X)}{P(X)}$; D) $\frac{P(A \cup X)}{P(A)}$;
 E) $\frac{P(A \cap X)}{P(A)}$; F) $\frac{P(X)}{P(A \cap X)}$.

60. Bernulli formulasi $P_{m,n} =$

- A) $C_m^n p^m q^{m-n}$; B) $C_m^n p^n q^{m-n}$; D) $C_m^n p^{m-n} q^n$;
 E) $C_n^m p^{n-m} q^m$; F) $C_n^m p^m q^{n-m}$.