

.....3.

I . $p -$ $p -$ 5.

1.1.5.

1.2. $p -$ 8.

1.3. $p -$ 10.

1.4. $p -$ 11.

II . $p -$ 12.

2.1.12.

2.2. $p -$ 14.

.....20.

.....21.

$p -$

$R -$

$C -$

p

F_p

$$|x + y| \leq \max(|x|, |y|)$$

$Q_p - p -$

K

J.C.Yoccoz [3]

$p -$

$p -$

$p -$

([1], [3]).

$$f : K \rightarrow K$$

1997

[3]

$p -$

Q_p

$p -$

$p -$

$p -$

$p -$

$p > 1$

$p -$

$Q_p p -$

[5]

C_p

$p -$

$$\left(x^n, \frac{x+a}{bx+c}, \frac{ax^2}{bx+1} \right)$$

$p -$

$$f(x) = \frac{x + a + xA(x)}{bx + c + A(x)}$$

$p -$

1.1. $\mathbf{I} \cdot p - \quad p - \cdot$
 $\dots(x, y)$, (x, y)

- 1) $\dots(x, y) \geq 0 \quad \dots(x, y) = 0 \Leftrightarrow x = y$,
 2) $\dots(x, y) = \dots(y, x), \quad \forall x, y \in X$
 3) $\forall z \in X \quad \dots(x, y) \leq \dots(x, z) + \dots(z, y), \quad (\quad)$,
 $\dots(x, y) \quad (X, \dots)$

$$\dots(x, y) \leq \max\{\dots(x, z), \dots(z, y)\} \quad (\quad) \quad (1.1)$$

(X, \dots)

1.1.

$$\dots(a, c) = \max\{\dots(a, b), \dots(b, c)\} \quad a, b, c \in X \quad \dots(a, b) \neq \dots(b, c),$$

$$\dots(a, b) < \dots(b, c)$$

$$\dots(a, c) \leq \max\{\dots(a, b), \dots(b, c)\} = \dots(b, c)$$

$$\dots(a, b) < \dots(b, c). \quad \dots(b, c) \leq \max\{\dots(a, b), \dots(a, c)\} = \dots(a, c)$$

$$\dots(a, c) = \max\{\dots(a, b), \dots(b, c)\}.$$

1.2. (X, \dots)

$$U_r(a) = \{x \in X : \dots(a, x) < r\}$$

$$a \in X \quad r \in \mathbb{R}^+$$

$$V_r(a) = \{x \in X : \dots(a, x) \leq r\}$$

$$a \quad r$$

$$S_r(a) = \{x \in X : \dots(a, x) = r\}$$

$$a \quad r$$

1.3. (X, \dots)

$$V_r(a)$$

$$b \in V_r(a)$$

$$V_r(b) = V_r(a)$$

$$x \in V_r(b)$$

$$\dots(x, a) \leq \max\{\dots(x, b), \dots(b, a)\} \leq r.$$

$$V_r(b) \subseteq V_r(a).$$

$$V_r(a) \subseteq V_r(b).$$

$$V_r(b) = V_r(a).$$

1.4. $a, b \in X \quad r, s \in \mathbb{R}^+, \quad (0 < s \leq r)$

$$V_r(a) \quad V_s(b)$$

$$V_s(b) \subseteq V_r(a)$$

$$\begin{aligned}
 & F, \|\cdot\|_1, \|\cdot\|_2 \\
 & 1.2. p - P, Q, n, m \\
 & \forall x \in Q \setminus \{0\} \\
 & x = p^{x(x)} \frac{m}{n}, \quad n \in N, \quad x = x(x), m \in Z, \quad (m, p) = (n, p) = 1 \quad (1.3)
 \end{aligned}$$

$$|x|_p = \begin{cases} p^{-x(x)}, & x \neq 0 \\ 0, & x = 0 \end{cases} \quad (1.4)$$

$$1.2.1. |\cdot|_p \quad Q$$

$$1) |x|_p \geq 0 \quad \forall x \in Q \quad |x|_p = 0 \Leftrightarrow x = 0.$$

$$2) |xy|_p = |x|_p |y|_p, \quad \forall x, y \in Q.$$

$$3) |x + y|_p \leq \max\{|x|_p, |y|_p\}, \quad \forall x, y \in Q$$

$$1) \quad 2) \quad 3)$$

$$\forall x, y \in Q \quad (3) \quad x = p^{x(x)} \frac{m_1}{n_1}, \quad y = p^{x(y)} \frac{m_2}{n_2}$$

$$\min\{x(x), x(y)\} = x(x)$$

$$x + y = p^{x(x)} \frac{m_1}{n_1} + p^{x(y)} \frac{m_2}{n_2} = p^{x(x)} \frac{m_1 n_2 + m_2 n_1 p^{x(y)-x(x)}}{n_1 n_2}$$

$$n_1 n_2 \quad p, \quad m_1 n_2 + m_2 n_1 p^{x(y)-x(x)} \quad p$$

$$x(x + y) \geq x(x) = \min\{x(x), x(y)\}$$

$$|x + y|_p = p^{-x(x+y)} \leq \max\{p^{-x(x)}, p^{-x(y)}\} = \max\{|x|_p, |y|_p\}$$

$$1.2.2. \quad |x|_p \neq |y|_p, \quad |x + y|_p = \max\{|x|_p, |y|_p\}$$

$$1.2.3. \quad p = 2, \quad |x|_2 = |y|_2, \quad |x + y|_2 \leq \frac{1}{2} |x|_2$$

$$1.2.4. \quad Q, |\cdot|_p, p -$$

Q

|\cdot|

R

Q

$$|\cdot|_p \quad p -$$

$$Q_p \quad p -$$

1.2.5. () [2] Q

$$p^{-1} \cdot p^{-1} \cdot \forall y \in R \setminus \{0\}$$

$$y = p^{x(y)}(y_0 + y_1 p^{-1} + y_2 p^{-2} + \dots),$$

$$y_j, x(y) \in Z, 0 \leq y_j \leq p-1, y_0 > 0, (j = 0, 1, 2, \dots).$$

$$, \forall x \in Q_p \setminus \{0\} p^{-1}$$

$$= p^{x(\cdot)}(y_0 + y_1 p^{-1} + y_2 p^{-2} + \dots),$$

$$y_j, x(\cdot) \in Z, 0 \leq y_j \leq p-1, y_0 > 0, (j = 0, 1, 2, \dots). \tag{1.5}$$

1.2.6. $x \in Q_p$ (1.5)

$$x_j, j = 0, 1, 2, \dots$$

$$\dots -1 = (p-1) + (p-1)p + (p-1)p^2 + \dots$$

1.3. p^{-1}
 Q_p

1.3.1. F $F[x]$

$$F \cdot F^{-1} \cdot F \cdot K$$

$$F = \overline{K}$$

$$R \cdot x^2 + 1 = 0 \cdot R$$

$$i, (i^2 = -1) \cdot R$$

$$Q_p \cdot Q_p[x] \cdot Q_p \cdot p^{-1} \cdot Q_p$$

$$Q_p(\cdot)$$

1.3.2. [2]. $Q_p(\cdot)$

$$Q_p(\cdot) \cdot p^{-1} \cdot C_p \cdot p^{-1}$$

1.3.3. () $C_p \cdot p^{-1}$

C_p [2]:

1^o. $\forall x \in C_p \exists r \in Q, |x|_p = p^r$

2^o. C_p

1.4. p -
 C_p p -
 $\{x_j\} \subset C_p, j=1,2,3,\dots$ $x \in C_p$, $\lim_{j \rightarrow \infty} |x_j - x|_p = 0$
 $\{x_j\}$ x

1.4.1. $D \subset C_p$ $x \in D$ $\forall v > 0$
 $\exists u > 0$, $|y - x|_p < u$ $\forall y \in D$ $|f(y) - f(x)|_p < v$
 $f : D \rightarrow C_p$ $x \in D$

1.4.2. $D \subset C_p$ $f : D \rightarrow C_p$ f
 $x \in D$
 $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $f'(\cdot), D$, f D

1.4.3. [3] C_p $\{a_n\}$
 $\sum_{n=0}^{\infty} a_n$ $\lim_{n \rightarrow \infty} |a_n|_p = 0$
 $f : C_p \rightarrow C_p$ p -

$$f(x) = \sum_{n=0}^{\infty} f_n x^n, f_n \in C_p \quad (1.6)$$

1.4.4. (1.6) $|x|_p \leq R$ $|x|_p > R$
 $R = R(f)$, (1.6)

$$R = \frac{1}{\lim_{n \rightarrow \infty} |f_n|_p^{1/n}} \quad (1.7)$$

1.4.5. $U_r(a) \subset C_p$ $f : U_r(a) \rightarrow C_p$

$$f(x) = \sum_{n=0}^{\infty} f_n (x-a)^n, f_n \in C_p \quad (1.8)$$

$f(x)$ $U_r(a)$

II p -

$$D \subset C_p \quad f : D \rightarrow C_p \quad (f, D)$$

$$\forall x_0 \in D \quad x_n = f^n(x_0), \quad n = 1, 2, 3, \dots$$

$$f^n(x_0) = \underbrace{f \circ f \circ \dots \circ f}_n(x_0).$$

2.1.

$$(f, D) \quad 2.1.1. \quad x_0 \in D \quad f(x_0) = x_0, \quad x_0$$

$$2.1.2. \quad x_0 \in D \quad U_r(x_0) \quad \lim_{n \rightarrow \infty} f^n(x) = x_0, \quad x_0$$

$$A(x_0) = \{ x \in D : \lim_{n \rightarrow \infty} f^n(x) = x_0 \}$$

$$2.1.3. \quad x_0 \in D \quad U_r(x_0) \quad \forall x \in U_r(x_0) \quad |x - x_0|_p < |f(x) - x_0|_p, \quad x_0$$

$$2.1.4. \quad x_0 \in D, \quad f(x) \in D \quad U_r(x_0) \quad \forall \dots < r \quad S_{\dots}(x_0) \quad f(x) \in U_r(x_0) \quad x_0$$

$$D \quad x_0$$

$$SI(x_0)$$

$$2.1.5. \quad x_0 \in D, \quad f(x) \in D \quad |f'(x_0)|_p < 1, \quad x_0 \quad ; \quad |f'(x_0)|_p = 1, \quad x_0$$

$$|f'(x_0)|_p > 1, \quad x_0$$

$$2.1.6. \quad x_0 \in D, \quad f(x) \in D$$

$$1^0. \quad x_0, \quad f(x) \in D \quad r > 0$$

$$q = \max_{1 \leq n < \infty} \left| \frac{1}{n!} \frac{d^n f}{dx^n}(x_0) \right|_p r^{n-1} < 1 \quad (2.1)$$

$$2^0. \quad U_r(x_0) \subset D, \quad U_r(x_0) \subset A(x_0)$$

$$x_0, \quad f(x) \in D, \quad r > 0$$

$$s = \max_{2 \leq n < \infty} \left| \frac{1}{n!} \frac{d^n f}{dx^n}(x_0) \right|_p r^{n-1} < |f'(x_0)|_p \quad (2.2)$$

$$3^0. \quad U_r(x_0) \subset D, \quad U_r(x_0) \subset SI(x_0), \quad (f, D)$$

$$2.1.7. [1] \left(\begin{array}{l} f(x) \\ V_r(x_0), \dots \leq r \end{array} \right), \quad \left(\begin{array}{l} f'(x_0) \neq 0, |f'(x_0)|_p = p^n \\ V_{\dots+n}(y_0), y_0 = f(x_0) \\ g(y) \\ V_{\dots+n}(y_0) \end{array} \right)$$

$$g'(y_0) = \frac{1}{f'(x_0)}; \quad |g(y) - x_0|_p = |g'(y_0)|_p |y - y_0|_p, \quad y \in V_{\dots+n}. \quad (2.3)$$

2.2. $p -$

$$D \subset C_p \quad f: D \rightarrow C_p$$

$$f(x) = \frac{x + a + xA(x)}{bx + c + A(x)}, \quad b \neq 0, \quad a, b, c \in C_p \quad (2.4)$$

$$A(x) \neq -bx - c.$$

(2.4)

$$f(x) = x, \quad x_{1,2} = \frac{1 - c \pm \sqrt{(c-1)^2 + 4ab}}{2b} \quad (2.5)$$

$$A(x) \quad D$$

$$f'(x) = \frac{c - ab + (1+c)A(x) + (bx^2 + (c-1)x - a)A'(x) + A^2(x)}{(bx + c + A(x))^2}$$

(2.5)

$$f'(x_j) = \frac{c - ab + (1+c)A'(x_j) + A^2(x_j)}{(bx_j + c + A(x_j))^2}, \quad j = 1, 2. \quad (2.6)$$

$$t^2 = \frac{(c-1)^2 + 4ab}{4},$$

$$n_1 = 2A(x_2) + 1 + c + t,$$

$$n_2 = 2A^2(x_2) + 2tA(x_2) + 4ab - 4c - t,$$

$$n_3 = (1+c-t)A^2(x_2) - (2ab - 2c - t)A(x_2),$$

2.2.1.

$$A(x_1) = \mathbb{E}(A(x_2)), \quad A(x_1) = \{A(x_2)\}$$

$$1. \quad x_1 \quad (x_2) \quad , \quad x_2 \quad (x_1)$$

$$2. \quad x_1 \quad , \quad x_2$$

$$\{ (A(x_2)) = \frac{u_2 + \sqrt{u_2^2 + 4u_1u_3}}{2u_1}, \quad \mathbb{E}(A(x_2)) = \frac{u_2 - \sqrt{u_2^2 + 4u_1u_3}}{2u_1}$$

$$f'(x_1)f'(x_2) = 1 \quad (2.7)$$

$$(2.8) \quad u_1 A^2(x_1) - u_2 A(x_1) - u_3 = 0 \quad (2.7)$$

$$\begin{aligned} & [c - ab + (1 + c)A(x_1) + A^2(x_1)][(1 + c - 2t) + 2A(x_2)]^2 = \\ & = [c - ab + (1 + c)A(x_2) + A^2(x_2)][(1 + c + 2t) + 2A(x_1)]^2 \end{aligned}$$

$$A(x) = x$$

$$f(x) = \frac{x^2 + x + a}{bx + c}, \quad b \neq 0, \quad a, b, c \in C_p \quad (2.9)$$

$$f(x)$$

$$1- \quad b = 1, \quad c \neq 1, \quad x_0 = \frac{a}{c-1};$$

$$2- \quad b \neq 1, \quad c \neq 1, \quad x_{1,2} = \frac{1-c \pm \sqrt{(c-1)^2 + 4a(b-1)}}{2(b-1)};$$

$$3- \quad b = 1, \quad c = 1, \quad a \neq 0$$

$$1- \quad (2.9)$$

$$f(x) = \frac{x^2 + x + a}{x + c} \quad (2.10)$$

$$x \neq \tilde{x} = -c.$$

$$\Pi = \{x \in C_p : \exists n \in N, f^n(x) = \tilde{x}\} \quad \Pi \quad \tilde{x}$$

$$(f, C_p \setminus \Pi)$$

$$(2.10)$$

$$f'(x) = \frac{x^2 + 2cx + c - a}{(x + c)^2},$$

$$f^{(n)}(x) = (-1)^n n! \frac{c^2 - c + a}{(x + c)^{n+1}}$$

$$x_0 = \frac{a}{c-1}$$

$$f'(x_0) = \frac{x_0^2 + 2cx_0 + c - a}{(x_0 + c)^2} = \frac{a + c - 1}{c^2 - c + a} \quad (2.11)$$

$$f(y) \in S_-(x_0), \quad y \quad |f(y) - x_0|_p = \sim, \\ f(S_-(x_0)) \subset S_-(x_0).$$

$$2.2.4. \quad |f'(x_0)|_p < 1, \quad U_u(x_0) \subset A(x_0), \\ 2.1.6, \quad (2.15) \quad r > 0$$

$$q = \max_{1 \leq n < \infty} \left| \frac{1}{n!} \frac{d^n f}{dx^n}(x_0) \right|_p r^{n-1} < 1 \quad (2.17)$$

$$V_r(x_0) \subset C_p, \quad V_r(x_0) \subset A(x_0)$$

$$q = \max_{1 \leq n < \infty} \left| \frac{1}{n!} \frac{d^n f}{dx^n}(x_0) \right|_p r^{n-1} = \max_{1 \leq n < \infty} \left| \frac{c^2 - c + a}{(x_0 + c)^{n+1}} \right|_p r^{n-1} = \\ = \max_{1 \leq n < \infty} \left| \frac{c^2 - c + a}{(x_0 + c)^2} \right|_p \left| \frac{1}{(x_0 + c)^{n-1}} \right|_p r^{n-1} = \max_{1 \leq n < \infty} \left| \frac{(c-1)^2}{c^2 - c + a} \right|_p \left| \frac{1}{x_0 + c} \right|_p^{n-1} r^{n-1} = \\ = \max_{1 \leq n < \infty} \left| \frac{c^2 - c + a - a - c + 1}{c^2 - c + a} \right|_p \frac{r^{n-1}}{|x_0 + c|_p^{n-1}} =$$

$$= \max_{1 \leq n < \infty} |1 - f'(x_0)|_p \frac{r^{n-1}}{|x_0 + c|_p^{n-1}} \leq \max_{1 \leq n < \infty} \left| \frac{r}{x_0 + c} \right|_p^{n-1}.$$

$$r < u = |x_0 + c|_p, \quad (2.17)$$

$$U_u(x_0) \subset A(x_0).$$

$$2.2.5. \quad \Pi \subset S_u(x_0).$$

$$\tilde{x} \in \Pi \quad S_u(x_0),$$

$$\tilde{x} = -c, \quad |-c - x_0|_p = |x_0 + c|_p = u. \quad 2.2.3. \quad 2.2.4$$

$$\Pi \subset S_u(x_0).$$

$$x \in S_u(x_0) \setminus \Pi$$

$$2.2.6. \quad |f'(x_0)|_p < 1 \quad x \in S_u(x_0) \setminus \Pi$$

$$f^k(x) \in S_{-k}(x_0), \quad k \in N, \quad \sim_k > u.$$

$$x \in S_u(x_0) \setminus \Pi, \quad |x - x_0|_p = u.$$

$$x = x_0 + \langle, \quad |\langle|_p = u \quad (2.14)$$

$$|f(x) - x_0|_p = \frac{\left| \left\langle \left| f'(x_0) + \frac{\langle}{x_0 + c} \right| \right\rangle \right|_p}{\left| 1 + \frac{\langle}{x_0 + c} \right|_p} \geq u.$$

$$\begin{aligned} |f(x) - x_0|_p > u & , & \sim_1 > u & , \\ f(x) \in S_{-1}(x_0) & . & 2.2.3 & . \\ |f(x) - x_0|_p = u & , & & \end{aligned}$$

$$|f^2(x) - x_0|_p = \frac{|f(x) - x_0|_p \left| f'(x_0) + \frac{f(x) - x_0}{x_0 + c} \right|_p}{\left| 1 + \frac{f(x) - x_0}{x_0 + c} \right|_p} = \frac{u^2}{|f(x) + c|_p}.$$

$$f(x) \in S_u(x_0) \quad f(x) = x_0 + \langle', \left| \langle' \right|_p = u .$$

$$|f^2(x) - x_0|_p = \frac{u^2}{|f(x) + c|_p} = \frac{u^2}{\left| \langle' + (x_0 + c) \right|_p} .$$

$$, |f^2(x) - x_0|_p \geq u .$$

$$|f^2(x) - x_0|_p > u \quad , \quad \sim_2 > u \quad , \quad f^2(x) \in S_{-2}(x_0)$$

$$|f^2(x) - x_0|_p = u \quad ,$$

$$|f^{k-1}(x) - x_0|_p = u \quad , \quad (2.14)$$

$$|f^k(x) - x_0|_p = \frac{|f^{k-1}(x) - x_0|_p \left| f'(x_0) + \frac{f^{k-1}(x) - x_0}{x_0 + c} \right|_p}{\left| 1 + \frac{f^{k-1}(x) - x_0}{x_0 + c} \right|_p} = \frac{u^2}{|f^{k-1}(x) + c|_p}.$$

$$f^{k-1}(x) \in S_u(x_0) \quad f^{k-1}(x) = x_0 + \langle''', \left| \langle''' \right|_p = u .$$

$$|f^k(x) - x_0|_p = \frac{u^2}{|f^{k-1}(x) + c|_p} = \frac{u^2}{\left| \langle''' + (x_0 + c) \right|_p} .$$

$$, |f^k(x) - x_0|_p \geq u . \quad |f^k(x) - x_0|_p > u \quad , \quad \sim_k > u$$

$$, f^k(x) \in S_{-k}(x_0) .$$

$$|f^k(x) - x_0|_p = u \quad ,$$

